



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

This is the equation of the curve described by the dog, and it is called the "Curve of pursuit." When the dog overtakes the hare, $y=0$, $x=ON=b$.

\therefore (12) becomes, $\frac{na}{n^2-1}=b\dots(13)$. Solving (13) as a quadratic in n , we

have, $n=\frac{1}{2b}(a\pm\sqrt{4b^2+a^2})$, or $n=1.0644+$. Substituting this value of n in

(2), remembering that when the dog overtakes the hare, $EP=s$, $x=320$, $y=0$, we have, $s=1.0644\times 320=340.624+$ rods.

PROBLEMS.

33. Proposed by Professor ALEXANDER ROSS, C. E., Sebastopol, California.

From a point P without a rectangular field $ABCD$, the distances PA , PB , and PC measured to the corners are, respectively, 70, 40, and 60 chains. What is the area of the field?

34. Proposed by THOS. U. TAYLOR, C. E., M. C. E., Department of Engineering, University of Texas, Austin, Texas.

Given a variable parallelogram $ABCP$, where P remains fixed. A moves on an irregular plane curve (closed) and C moves on another plane curve (closed) whose plane is parallel to the plane of (A) curve. The generator PC moves completely around and returns to its initial position, AB always moving parallel to PC , and, of course, returns to its initial position. If distances between planes (A) and ($C=h$), show by elementary mathematics and without using theorem of Koppé that volume of solid generated by variable parallelogram $ABCP=\frac{1}{2}h$ (area generated by AP + area generated by BC).

QUERIES AND INFORMATION.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

SPACE.

Space is an entity, outside of the human mind, extended in three directions at right angles to each other, continues, immaterial, immovable, inflexible and illimitable. It is an entity, sui generis, neither psychical nor physical.

It is cognized but not created by the mind of man and is, doubtless, what it is cognized to be.

A fourth dimension has never been discovered.

Arthur Willink in "The World of the Unseen," pages 90 and 91, locates his hypothetical "Higher Space" in an unknown direction from our space.